

IMAGING SERVICES OUT-PATIENT BOOKING

100 mm 10	100 100 100 100 100 100 100 100 100 100	AP	POINTMENT					**************************************	**************************************
DATE:	MTWTHF	s s	TIME:		вос	OKED	BY:		·
			HARGED IF A MINI SED OR CANCEL				TS		
REQUIRED PATIENT INFORMATION									
PATIENT'S SURNAME:	3. Miles 2 d. m		GIVEN NAMES:			der ein de			
BIRTH DATE:	057 14 5		MAIDEN NAME:					 	
YY MM DD	SEX M	F 🗆						···	
PHN and/or MRN:									
ADDITIONAL PATIENT INFORMATION									
ADDRESS			Phone No.(H)	_					
			Phone No.(W)						
Postal Code:			Financial Class	MSP □	WC	В□	Self	Pay 🗆 O	her 🗆
FOR RAD DAY CARE PATIENTS DCR D DCO D			WCB #:						
	CLIN	ICAL	INFORMATIO	N					-
Procedure/Type of Exam			Previous X-rays						
Tentative Diagnosis									
History and Clinical Diagnosis including date onset of symptoms									
Ordering Doctor:	rdering Doctor: MSP			# Signature:					
Copy of Report to: (I)			(2)						
Previous Examination results:					_				
X-rays Head			Chest Abdomen Pelvis Spine Other						
C.T.									
Ultrasound									
Nuclear Medicine									
ECG								Without	With
Previous back surgery Yes □	No 🗆			Telebrix	(
Allergies Yes ☐ No ☐				Water					
If yes (specify) Diabetic Yes □ No □				Intraver	nous (Contr	ast		
If yes, is the patient taking Gluoph	age			Slice W	idth				
(Metformin) Yes □ No □				Priority		L	Appr Time	oximate	

Guidelines for Ordering CT Scan in Adults with Acute Head Injury

Absolute indications:

- 1) GCS < 15 in the absence of intoxicant
- 2) Deteriorating level of consciousness
- 3) Depressed skull fracture

Consider CT scan in the following situation:

- 1) GCS 15 At the discretion of the physician. Risk factors to consider:
 - Abnormal examination suggesting a focal lesion
 - Skull fracture in children
 - Extremes of age
 - Mechanism of injury such as pedestrian struck
 - Bleeding diathesis
 - Severe persistent symptoms such as headache or vomiting
 - Prolonged loss of consciousness
 - Multiple trauma with head injury requiring general anesthetic
- 2) GCS < 15 with intoxicant Consider observation initially. CT scan at discretion of physician if:
 - Patient has other risk factors
 - Level of consciousness does not improve or is inconsistent with blood alcohol level

A.1133NS Side 2